CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, ASCA, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal and Science Citation Index)

Prediction of retention in gas–liquid chromatography using the UNIFAC group contribution method. I. Low-molecular-weight stationary phases
by G. J. Price (Bath, U.K.) and M. R. Dent (London, U.K.) (Received August 14th, 1989) 1

Gas–liquid chromatography–mass spectrometry of mono- and dithiols as their *tert.*-butyldimethylsilyl derivatives
by D. C. Landrum and T. P. Mawhinney (Columbia, MO, U.S.A.) (Received August 28th, 1989) 21

Evaluation of 4-(2′-phenylisopropyl)phenol esters as stationary phases for gas–liquid chromatography
by N. R. Ayyangar, A. S. Tambe and S. S. Biswas (Pune, India) (Received August 23rd, 1989) 33

Capillary gas chromatographic determination of volatiles in solid matrices by direct introduction using a programmable-temperature vaporizer
by G. Reglero, M. Herraiz, T. Herraiz and J. Sanz (Madrid, Spain) (Received August 8th, 1989) 43

Packed-column supercritical fluid chromatography of benzodiazepines
by R. M. Smith and M. M. Sanagi (Loughborough, U.K.) (Received August 11th, 1989) . 51

Preparative high-performance liquid chromatography under isocratic conditions. I. Craig simulations for heavily overloaded separations
by L. R. Snyder and J. W. Dolan (Lafayette, CA, U.S.A.) and G. B. Cox (Glasgow, DE, U.S.A.) (Received August 15th, 1989) 63

Preparative high-performance liquid chromatography under isocratic conditions. II. The role of column variables
by L. R. Snyder (Lafayette, CA, U.S.A.) and G. B. Cox (Glasgow, DE, U.S.A.) (Received August 15th, 1989) 85

Preparative high-performance liquid chromatography under isocratic conditions. III. The consequences of two adjacent bands having unequal column capacities
by G. B. Cox (Glasgow, DE, U.S.A.) and L. R. Snyder (Lafayette, CA, U.S.A.) (Received August 15th, 1989) 95

Automatic on-line pressurized liquid injection in gas chromatography
by J. H. Marsman, H. J. Panneman and A. A. C. M. Beenackers (Groningen, The Netherlands) (Received September 5th, 1989) 111

Rapid determination of seven herbicides in water or isooctane using C_{18} and Florisil Sep-Pak cartridges and gas chromatography with electron-capture detection
by W.-K. Wang and S.-D. Huang (Hsinchu, Taiwan) (Received August 28th, 1989) . . . 121

Synthetic peptide purification by application of linear solvent strength gradient theory
by J. C. Ford and J. A. Smith (Boston, MA, U.S.A.) (Received September 19th, 1989) . . 131

Separation of inorganic anions by ion chromatography on dye-coated silica gel RP-18
by G. Müller and H.-U. Meisch (Saarbrücken-Dudweiler, F.R.G.) (Received September 4th, 1989) 145

(Continued overleaf)

Contents (continued)

Gradient high-performance liquid chromatography using alkylphenone retention indices of insecticidal extracts of *Penicillium* strains
by R. R. M. Paterson (Kew, U.K.) and C. Kemmelmeier (Maringa, Brazil) (Received August 8th, 1989) 153

Separation of homologous and isomeric alkaloids related to nicotine on a β-cyclodextrin-bonded phase
by J. I. Seeman and H. V. Secor (Richmond, VA, U.S.A.) and D. W. Armstrong, K. D. Ward and T. J. Ward (Rolla, MO, U.S.A.) (Received August 16th, 1989) 169

Theory of isotope separation by displacement chromatography
by H. Kakihana and T. Oi (Tokyo, Japan) (Received September 1st, 1989) 179

Effect of organic solvents in sample solutions and injection volumes on chromatographic peak profiles of analytes in reversed-phase high-performance liquid chromatography
by D. Vukmanic and M. Chiba (St. Catharines, Canada) (Received August 22nd, 1989) . 189

Preparation of carboneous adsorbents by catalytic decomposition of hydrocarbons for chromatographic application
by A. Gierak and R. Leboda (Lublin, Poland) (Received August 11th, 1989) 197

Spectroscopic investigation of the hydrolysis of γ-glycidoxypropylsilane bound to silica surfaces
by G. R. Bogart and D. E. Leyden (Fort Collins, CO, U.S.A.) and T. M. Wade, W. Schafer and P. W. Carr (Minneapolis, MN, U.S.A.) (Received September 18th, 1989) 209

Determination of diffusion coefficients of proteins in beaded agarose by gel filtration
by P. A. Davies (Oxford, U.K.) (Received July 26th, 1989) 221

Study of the chromatographic behaviour and resolution of α-amino acid enantiomers by high-performance liquid chromatography utilizing pre-column derivatization with *o*-phthaldialdehyde and new chiral thiols
by M. R. Euerby, L. Z. Partridge and W. A. Gibbons (London, U.K.) (Received August 18th, 1989) 239

Reversed-phase high-performance liquid chromatographic purification of ^{125}I-labeled rat growth hormone-releasing hormone for radioimmunoassay
by J. W. Leidy, Jr. (Huntington, WV, U.S.A.) (Received August 23rd, 1989) 253

Determination of ion-exchange equilibrium parameters of amino acid and protein systems by an impulse response technique
by R. D. Whitley, J. M. Brown, N. P. Karajgikar and N.-H. L. Wang (West Lafayette, IN, U.S.A.) (Received August 29th, 1989) 263

Interaction of lactate dehydrogenase with structurally related triazine dyes using affinity partitioning and affinity chromatography
by J. Kirchberger (Leipzig, G.D.R.), F. Cadelis (Compiègne, France), G. Kopperschläger (Leipzig, G.D.R.) and M. A. Vijayalakshmi (Compiègne, France) (Received June 14th, 1989) 289

Microheterogeneity of the growth-associated neuronal protein B-50. Contribution of phosphorylation by protein kinase C
by P. Schotman, H. B. Nielander, A. J. van Rozen, H. Frankena, L. H. Schrama and W. H. Gispen (Utrecht, The Netherlands) (Received July 11th, 1989) 301

Purification of tryptophan decarboxylase from a *Catharanthus roseus* cell suspension culture
by E. J. M. Pennings, R. Verpoorte, O. J. M. Goddijn and J. H. C. Hoge (Leiden, The Netherlands) (Received August 8th, 1989) 311

High-performance liquid chromatography determination of the extent of proteolysis in Gouda cheese
by W. Haasnoot, P. Stouten and D. P. Venema (Wageningen, The Netherlands) (Received June 22nd, 1989) 319

Preparative purification of functional bacteriorhodopsin by high-performance size-exclusion chromatography
by L. J. W. Miercke and R. M. Stroud (San Francisco, CA, U.S.A.) and E. A. Dratz (Bozeman, MT, U.S.A.) (Received August 15th, 1989) 331

High-performance liquid chromatographic determination of low levels of primary and secondary amines in aqueous media via derivatisation with 1,2-naphthoquinone-4-sulphonate
by J. R. L. Smith and A. U. Smart (York, U.K.) and F. E. Hancock and M. V. Twigg (Billingham, U.K.) (Received September 28th, 1989) 341

Proposed primary reference methods for the determination of some commercially important chiral aryloxypropionate herbicides in both free acid and ester forms
by B. Blessington and N. Crabb (Bradford, U.K.) (Received August 29th, 1989) 349

Centrifugal counter-current partition chromatography with helical coil rotor. Simplified counter-current chromatography with a rotating face-seal
by L.-S. Kim, Z.-Q. Tian, X.-N. Li, Y. Zhang and J. Wang (Beijing, China) (Received July 17th, 1989) 359

Improved separation of lipid esters by thin-layer chromatography
by H. K. Bhat and G. A. S. Ansari (Galveston, TX, U.S.A.) (Received August 15th, 1989) 369

Notes

Modification to a thermal energy analyzer with associated electronic filtering for improved gas chromatographic analysis of explosives traces
by D. A. Collins (Birmingham, U.K.) (Received August 29th, 1989) 379

Identification and analysis of tetramethylnaphthalenes in petroleum
by P. G. Forster, R. Alexander and R. I. Kagi (Perth, Australia) (Received July 20th, 1989) 384

Correlation between chromatography and ^{1}H NMR parameters in sterically substituted phenols
by S. R. Salman (Baghdad, Iraq), F. S. Kamounah (Basrah, Iraq) and A. H. H. Tameesh (Baghdad, Iraq) (Received July 4th, 1989) 390

Low-temperature side-chain liquid crystalline polysiloxanes used as stationary phases for capillary gas chromatography
by Y. Jin, R. Fu, Z. Guan, J. Gong and B. Li (Beijing, China) (Received August 8th, 1989) 394

High-performance hydrophobic interaction chromatography of proteins on a pellicular support based on hydrophilic resin
by Y. Kato, T. Kitamura, S. Nakatani and T. Hashimoto (Yamaguchi, Japan) (Received June 22nd, 1989) 401

High-performance gel permeation chromatography of meningococcal polysaccharides
by P. Newland, B. Bingham, E. Tarelli and A. H. Thomas (Potters Bar, U.K.) (Received July 3rd, 1989) 406

Regiochemical differences in the isotopic fractionation of deuterated benzoic acid isotopomers by reversed-phase high-performance liquid chromatography
by W. J. S. Lockley (Loughborough, U.K.) (Received August 29th, 1989) 413

Separation of ^{125}I-labelled prostanoid derivatives by reversed-phase high-performance liquid chromatography
by I. Mucha and G. Tóth (Budapest, Hungary) (Received July 10th, 1989) 419

Determination of cyclic nucleotide phosphodiesterase activity in cellular systems by ion-pair reversed-phase liquid chromatography
by D. Techel and T. Braumann (Bremen, F.R.G.) (Received September 5th, 1989) 427

Improved high-performance liquid chromatographic method for artifact-free measurements of aldehydes in the presence of ozone using 2,4-dinitrophenylhydrazine
by D. F. Smith, T. E. Kleindienst and E. E. Hudgens (Research Triangle Park, NC, U.S.A.) (Received September 14th, 1989) 431

(Continued overleaf)

Contents (continued)

Analysis of *o*-phthalaldehyde derivatives of acidic and polar amino acids in fermentation broths by high-performance liquid chromatography
by R. L. White, A. C. Demarco and K. C. Smith (Wolfville, Canada) (Received September 19th, 1989) 437

High-performance liquid chromatography of triangular ruthenium clusters containing bidentate group-15 donor ligands
by A. Casoli, A. Mangia and G. Predieri (Parma, Italy) and E. Sappa (Torino, Italy) (Received July 3rd, 1989) 443

High-performance liquid chromatographic determination of pyrroloquinoline quinone with electrochemical detection and its application to bacterial samples
by M. Otani, C. Umezawa and K. Sano (Kobe, Japan) (Received August 22nd, 1989) . . 448

Synthesis of 7α-thiohydroxycholesterol involving the resolution of epimeric thiocholesterols by high-performance liquid chromatography
by J. A. Williams, D. S. Davies and S. Murray (London, U.K.) (Received July 6th, 1989) 454

Separation of diastereomeric amino acid conjugates of jasmonic acid
by G. Schneider, R. Kramell and C. Brückner (Halle/Saale, G.D.R.) (Received August 17th, 1989) 459

Author Index 463

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)